

## VGLUT2 (SLC17A6)

Cat.No. 135 418; Recombinant Guinea pig antibody, 50 µg recombinant IgG (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg purified recombinant IgG, lyophilized. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 up to 1 : 5000 (AP staining) (see remarks) <b>IP:</b> not tested yet <b>ICC:</b> 1 : 500 <b>IHC:</b> 1 : 500 <b>IHC-P:</b> 1 : 1000 <b>ExM:</b> external data (see remarks)
Clone	Gp95E11
Subtype	IgG2 (κ light chain)
Immunogen	Synthetic peptide corresponding to residues near the carboxy terminus of rat VGLUT2 (UniProt Id: Q9JI12)
Reactivity	Reacts with: rat (Q9JI12), mouse (Q8BLE7). Other species not tested yet.
Specificity	K.O. validated
Matching control	135-4P
Remarks	This antibody is a chimeric antibody based on the monoclonal mouse antibody clone 95E11. The constant regions of the heavy and light chains have been replaced by Guinea pig specific sequences. Therefore, the antibody can be used with standard anti-Guinea pig secondary reagents. The antibody has been expressed in mammalian cells. The antibody is highly recommended as a marker for glutamatergic nerve terminals. <b>WB:</b> To avoid protein aggregation, do not heat samples for SDS-PAGE. <b>ExM:</b> This antibody has been successfully applied and published for this method by customers (see application-specific references).

### Background

The vesicular **glutamate transporter 2 VGLUT2**, also referred to as **DNPI** and **SLC17A6**, has a more restricted expression than the related VGLUT1. Like VGLUT1, it is both necessary and sufficient for uptake and storage of glutamate and thus comprises the sole determinant for a glutamatergic phenotype. Both VGLUTs are different from the plasma membrane transporters in that they are driven by a proton electrochemical gradient across the vesicle membrane. VGLUT1 and VGLUT2 show complementary expression patterns. Together, they are currently the best markers for glutamatergic nerve terminals and glutamatergic synapses.

For more information on protein expression pattern, please refer to the overview image in our SYSY Antibodies ATLAS.

### Selected References for 135 418

- Glutamate stress in the caudal nucleus tractus solitarii (NTS): Impact on respiratory function and synaptic signaling in an Alzheimer's disease model.  
Carter SKR, Tipton RK, Forman CM, Mosher AP, Ostrowski D, Ostrowski TD  
Experimental neurology (2025) 387: 115190. . **WB; tested species: rat**
- Dopamine transporter and synaptic vesicle sorting defects underlie auxilin-associated Parkinson's disease.  
Vidyadhara DJ, Somayaji M, Wade N, Yücel B, Zhao H, Shashaank N, Ribaldo J, Gupta J, Lam TT, Sames D, Greene LE, et al.  
Cell reports (2023) 423: 112231. . **IHC; tested species: mouse**
- Loss of Bassoon in telencephalic excitatory neurons stabilizes non-aversive memories and strengthens aversive associations in social and non-social contexts.  
Zoicas I, Sucu B, Kornhuber J, Fejtova A  
Neurobiology of disease (2026) 222: 107362. . **IHC; tested species: mouse**
- A postsynaptic GPR158-PLCXD2 complex controls spine apparatus abundance and dendritic spine maturation.  
Verpoort B, Amado L, Vandendyck J, Leysen E, Dascenco D, Vandembempt J, Lemmens I, Wauman J, Vennekens K, Escamilla-Ayala A, Freitas ACN, et al.  
Developmental cell (2025) : . . **IHC; tested species: mouse**
- Altered motor coordination, vocal communication, and cerebellar circuit connectivity in mice carrying a near-complete human chromosome 21.  
Stander R, Ayyappan N, Sikorski D, van der Heijden ME, Hsiao K  
Translational psychiatry (2025) 161: 14. . **IHC; tested species: mouse**
- Inhibitory circuit dysfunction as a potential contributor to cortical reorganization in Glioblastoma progression.  
Spalletti C, Scalera M, Mori E, Haddad S, Mainardi M, Cangi D, Pillai V, Parmigiani E, Landi S, Caleo M, Vannini E, et al.  
Neurobiology of disease (2025) 213: 106997. . **IHC; tested species: mouse**
- Behavioral Modulation and Molecular Definition of Wide-Field Vertical Cells in the Mouse Superior Colliculus.  
Relota XJ, Ford A, Savier EL  
The Journal of neuroscience : the official journal of the Society for Neuroscience (2025) 4516: . . **IHC; tested species: mouse**

### Selected General References

- Identification of differentiation-associated brain-specific phosphate transporter as a second vesicular glutamate transporter (VGLUT2).  
Takamori S et al. J. Neurosci. (2001) PubMed:11698620
- The expression of vesicular glutamate transporters defines two classes of excitatory synapse.  
Fremeau RT et al. Neuron (2001) PubMed:11502256

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

Access the online factsheet including applicable protocols at <https://sysy.com/product/135418> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C

### Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.