

Rudolf-Wissell-Str. 28a 37079 Göttingen, Germany

Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

# VGLUT2 (SLC17A6)

Cat.No. 135 418; Recombinant Guinea pig antibody, 50 µg recombinant IgG (lyophilized)

## **Data Sheet**

Reconstitution/ Storage	50 µg purified recombinant IgG, lyophilized. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1: 1000 up to 1: 5000 (AP staining) IP: not tested yet ICC: 1: 500 IHC: 1: 500 IHC-P: 1: 1000 ExM: external data (see remarks)
Clone	Gp95E11
Subtype	IgG2 (κ light chain)
Immunogen	Synthetic peptide corresponding to residues near the carboxy terminus of rat VGLUT2 (UniProt Id: Q9JI12)
Reactivity	Reacts with: rat (Q9JI12), mouse (Q8BLE7). Other species not tested yet.
Specificity	K.O. validated
Matching control	135-4P
Remarks	This antibody is a chimeric antibody based on the monoclonal mouse antibody clone 95E11. The constant regions of the heavy and light chains have been replaced by Guinea pig specific sequences. Therefore, the antibody can be used with standard anti-Guinea pig secondary reagents. The antibody has been expressed in mammalian cells.  The antibody is highly recommended as a marker for glutamatergic nerve terminals.  WB: To avoid protein aggregation, do not heat samples for SDS-PAGE.  ExM: This antibody has been successfully applied and published for this method by customers (see application-specific references).

TO BE USED IN VITRO / FOR RESEARCH ONLY NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

# Background

The vesicular glutamate transporter 2 VGLUT2, also referred to as DNPI and SLC17A6, has a more restricted expression than the related VGLUT1. Like VGLUT1, it is both necessary and sufficient for uptake and storage of glutamate and thus comprises the sole determinant for a glutamatergic phenotype. Both VGLUTs are different from the plasma membrane transporters in that they are driven by a proton electrochemical gradient across the vesicle membrane.

VGLUT1 and VGLUT2 show complementary expression patterns. Together, they are currently the best markers for glutamatergic nerve terminals and glutamatergic synapses.

For more information on protein expression pattern, please refer to the overview image in our SYSY Antibodies ATLAS.

## Selected References for 135 418

Glutamate stress in the caudal nucleus tractus solitarii (nTS): Impact on respiratory function and synaptic signaling in an Alzheimer's disease model.

Carter SKR, Tipton RK, Forman CM, Mosher AP, Ostrowski D, Ostrowski TD Experimental neurology (2025) 387: 115190. WB; tested species: rat

Dopamine transporter and synaptic vesicle sorting defects underlie auxilin-associated Parkinson's disease.

Vidyadhara DJ, Somayaji M, Wade N, Yücel B, Zhao H, Shashaank N, Ribaudo J, Gupta J, Lam TT, Sames D, Greene LE, et al. Cell reports (2023) 423: 112231. IHC; tested species: mouse

A postsynaptic GPR158-PLCXD2 complex controls spine apparatus abundance and dendritic spine maturation.

Verpoort B, Amado L, Vandensteen J, Leysen E, Dascenco D, Vandenbempt J, Lemmens I, Wauman J, Vennekens K, Escamilla-Ayala A, Freitas ACN, et al.

Developmental cell (2025):.. IHC; tested species: mouse

#### **Selected General References**

Identification of differentiation-associated brain-specific phosphate transporter as a second vesicular glutamate transporter (VGLUT2).

Takamori S et al. J. Neurosci. (2001) PubMed:11698620

The expression of vesicular glutamate transporters defines two classes of excitatory synapse. Fremeau RT et al. Neuron (2001) PubMed:11502256

Immunocytochemical localization of candidates for vesicular glutamate transporters in the rat cerebral cortex. Fujiyama F et al. J. Comp. Neurol. (2001) PubMed:11406819

Molecular cloning of a novel brain-type Na(+)-dependent inorganic phosphate cotransporter. Aihara Y et al. J. Neurochem. (2000) PubMed:10820226

Access the online factsheet including applicable protocols at <a href="https://sysy.com/product/135418">https://sysy.com/product/135418</a> or scan the QR-code.



# FAQ - How should I store my antibody?

# **Shipping Conditions**

 All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freezedried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

# Storage of Sealed Vials after Delivery

- Unlabeled and biotin-labeled antibodies and control proteins should be stored at 4°C before reconstitution. They must not be stored in the freezer when still lyophilized!
   Temperatures below zero may cause loss of performance.
- Fluorescence-labeled antibodies should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

# Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle
  between freezing and thawing (to reduce frost-build-up), which is exactly what should be
  avoided. For the same reason, antibody vials should be placed in an area of the freezer that
  has minimal temperature fluctuations, for instance towards the back rather than on a door
  shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl)
  and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock
  concentration is affected by evaporation and adsorption of the antibody to the surface of the
  storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of
  activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

# **Product Specific Hints for Storage**

# Control proteins / peptides

• Store at -20°C to -80°C.

#### **Monoclonal Antibodies**

- Ascites and hybridoma supernatant should be stored at -20°C up to -80°C. Prolonged storage at 4°C is not recommended! Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### **Polyclonal Antibodies**

- Crude antisera: With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- Affinity purified antibodies: Less robust than antisera. Storage at -20°C up to -80°C is
  recommended. Adding a carrier protein like BSA will increase long term stability. Most of our
  antibodies already contain carrier proteins. Please refer to the data-sheet for detailed
  information.

#### Fluorescence-labeled Antibodies

• Store as a liquid with 1:1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

# FAQ - How should I reconstitute my antibody?

#### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add
  the amount of deionized water given in the respective datasheet. If higher volumes are
  preferred, add water as mentioned above and then the desired amount of PBS and a
  stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies
  already contain albumin. Take this into account when adding more carrier protein.
   For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the
  solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled
  with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1:1 (v/v) glycerol to a final
  concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in
  liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freezethaw cycles.
- Please refer to our tips and hints for subsequent storage of reconstituted antibodies and control peptides and proteins.