

Bassoon

Cat.No. 141 011; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

Data Sheet

| | |
|----------------------------|---|
| Reconstitution/ Storage | 100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For reconstitution add 100 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet. |
| Applications | WB: 1 : 1000 (AP staining) (see remarks) IP: external data (see remarks) ICC: 1 : 100 up to 1 : 500 (see remarks) IHC: not recommended IHC-P (FFPE): 1 : 500 ExM: external data (see remarks) ELISA: yes (see remarks) |
| Clone | 219E1 |
| Subtype | IgG2b (κ light chain) |
| Immunogen | Recombinant protein corresponding to residues near the carboxy terminus of rat Bassoon. (UniProt Id: O88778) |
| Reactivity | Reacts with: rat (O88778), mouse (O88737). Other species not tested yet. |
| Specificity | Specific for Bassoon |
| Matching control | 141-0P |
| Remarks | WB: Due to the large size of this protein, we recommend NuPAGE 3-8% Tris-Acetate gels for SDS-PAGE. IP: This antibody has been successfully applied and published for this method by customers (see application-specific references). ICC: 2% formaldehyde/PFA fixation is recommended. ExM: This antibody has been successfully applied and published for this method by customers (see application-specific references). ELISA: The ELISA-protocol for membrane proteins is required. Suitable as capture antibody for sandwich-ELISA. Please refer to the protocol for suitable detector antibodies. |

Background

Bassoon is a large protein which consists of an N-terminal Zn²⁺ finger and several piccolo-bassoon homology domains (PBH-domains). It is generally found together with piccolo, a related huge multi-domain protein of the CAZ (cytoskeletal matrix assembled at active zones). Bassoon was suggested to be a scaffolding element of the presynapse but deletion experiments in mice have shown that bassoon is also involved in synaptic vesicle cycling. Probably bassoon interacts with other protein factors via its Zn²⁺ domain but the potential partners have not been determined yet.

Selected References for 141 011

- Postsynaptic Y654 dephosphorylation of β-catenin modulates presynaptic vesicle turnover through increased n-cadherin-mediated transsynaptic signaling.
Chen CY, Chen YT, Wang JY, Huang YS, Tai CY
Developmental neurobiology (2017) 771: 61-74. . **WB, IP**
- Light-microscopy-based connectomic reconstruction of mammalian brain tissue.
Tavakoli MR, Lyudchik J, Januszewski M, Vistounou V, Agudelo Dueñas N, Vorlaufer J, Sommer C, Kreuzinger C, Oliveira B, Cenameri A, Novarino G, et al.
Nature (2025) 6428067: 398-410. . **EXM; tested species: mouse**
- eIF4B phosphorylation at Ser504 links synaptic activity with protein translation in physiology and pathology.
Bettgazzi B, Bellani S, Roncon P, Guarnieri FC, Bertero A, Codazzi F, Valtorta F, Simonato M, Grohovaz F, Zacchetti D
Scientific reports (2017) 71: 10563. . **ICC; tested species: rat**
- Remodelling of spared proprioceptive circuit involving a small number of neurons supports functional recovery.
Hollis ER, Ishiko N, Pessian M, Tolentino K, Lee-Kubli CA, Calcutt NA, Zou Y
Nature communications (2015) 6: 6079. . **IHC**
- Munc13-1 restoration mitigates presynaptic pathology in spinal muscular atrophy.
Moradi M, Weingart J, Deng C, Nasouti M, Briese M, Jablonka S, Sauer M, Sendtner M
Nature communications (2025) 161: 8724. . **ICC; tested species: mouse**
- Loss of synaptic Munc13-1 underlies neurotransmission abnormalities in spinal muscular atrophy.
Moradi M, Deng C, Sendtner M
Cellular and molecular life sciences : CMLS (2025) 821: 325. . **ICC; tested species: mouse**
- Input-Specific Localization of NMDA Receptor GluN2 Subunits in Thalamocortical Neurons.
Topolski MA, Gilmore BL, Khondaker R, Michniak JA, Studtmann C, Chen Y, Wagner GN, Pozo-Aranda AE, Farris S, Swanger SA
Journal of neurochemistry (2025) 1693: e70049. . **IHC; tested species: mouse**
- Real-time visualization of structural dynamics of synapses in live cells in vivo.
Son S, Nagahama K, Lee J, Jung K, Kwak C, Kim J, Noh YW, Kim E, Lee S, Kwon HB, Heo WD, et al.
Nature methods (2024) : . . **ICC; tested species: rat**
- Membrane compression by synaptic vesicle exocytosis triggers ultrafast endocytosis.
Ogunmowo TH, Jing H, Raychaudhuri S, Kusick GF, Imoto Y, Li S, Itoh K, Ma Y, Jafri H, Dalva MB, Chapman ER, et al.
Nature communications (2023) 141: 2888. . **ICC; tested species: mouse**
- Plasticity-Related Gene 5 Is Expressed in a Late Phase of Neurodifferentiation After Neuronal Cell-Fate Determination.
Gross I, Brandt N, Vonk D, Köper F, Wöhlbrand L, Rabus R, Witt M, Heep A, Plösch T, Hipp MS, Bräuer AU, et al.
Frontiers in cellular neuroscience (2022) 16: 797588. . **ICC; tested species: mouse**
- Dynamin is primed at endocytic sites for ultrafast endocytosis.
Imoto Y, Raychaudhuri S, Ma Y, Fenske P, Sandoval E, Itoh K, Blumrich EM, Matsubayashi HT, Mamer L, Zarebidaki F, Söhl-Kielczynski B, et al.
Neuron (2022) 11017: 2815-2835.e13. . **ICC; tested species: mouse**

Access the online factsheet including applicable protocols at <https://sysy.com/product/141011> or scan the QR-code.



TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.