

MARCO mouse-specific

Cat.No. HS-499 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 250 up to 1 : 500 (AP staining) IP: not tested yet ICC: not tested yet IHC: 1 : 500 (see remarks) IHC_P: 1 : 400 up to 1 : 1000
Immunogen	Synthetic peptide corresponding to residues near the carboxy terminus of mouse MARCO (UniProt Id: Q60754)
Reactivity	Reacts with: mouse (Q60754). No signal: rat (M0R9F7), human (Q9UEW3). Other species not tested yet.
Remarks	IHC: Heat-mediated antigen retrieval (in citrate buffer pH 6) is required for immunohistochemical staining

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

The macrophage receptor with collagenous structure (MARCO) belongs to the class A scavenger receptor (SR-A) family, all of which are transmembrane proteins with a homotrimeric structure (1). MARCO is constitutively expressed by peritoneal macrophages and splenic marginal zone macrophages. Its expression can be induced in other macrophages, such as Kupffer cells or alveolar macrophages, after bacterial infection or intravenous injection of LPS (2). MARCO mediates the binding of pathogens and unopsonized environmental particles, suggesting an important role in host defense (3). In tumor-associated macrophages, MARCO defines a subtype of suppressive TAMs (4) and is considered a potential prognostic and immunotherapeutic biomarker (5).

Selected General References

Scavenger receptor-A (CD204): a two-edged sword in health and disease.
Kelley JL, Ozment TR, Li C, Schweitzer JB, Williams DL
Critical reviews in immunology (2014) 343: 241-61. .

MARCO is a potential prognostic and immunotherapy biomarker.
Dong Q, Zhang S, Zhang H, Sun J, Lu J, Wang G, Wang X
International immunopharmacology (2023) 116: 109783. .

Reprogramming Tumor-Associated Macrophages by Antibody Targeting Inhibits Cancer Progression and Metastasis.
Georgoudaki AM, Prokopec KE, Boura VF, Hellqvist E, Sohn S, Östling J, Dahan R, Harris RA, Rantalainen M, Klevebring D, Sund M, et al.
Cell reports (2016) 159: 2000-11. .

Roles of a macrophage receptor with collagenous structure (MARCO) in host defense and heterogeneity of splenic marginal zone macrophages.
Ito S, Naito M, Kobayashi Y, Takatsuka H, Jiang S, Usuda H, Umezumi H, Hasegawa G, Arakawa M, Shultz LD, Elomaa O, et al.
Archives of histology and cytology (1999) 621: 83-95. .

Role of the scavenger receptor MARCO in alveolar macrophage binding of unopsonized environmental particles.
Palecanda A, Paulauskis J, Al-Mutairi E, Imrich A, Qin G, Suzuki H, Kodama T, Tryggvason K, Koziel H, Kobzik L
The Journal of experimental medicine (1999) 1899: 1497-506. .

Access the online factsheet including applicable protocols at <https://susy-histosure.com/product/HS-499003> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.