

## MafA

Cat.No. HS-521 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> not tested yet <b>IP:</b> not tested yet <b>IHC:</b> 1 : 1000 <b>IHC-P:</b> 1 : 2000 up to 1 : 4000 <b>ELISA:</b> not tested yet
Immunogen	Synthetic peptide corresponding to residues near the carboxy terminus of mouse MafA (UniProt Id: Q8CF90)
Reactivity	Reacts with: mouse (Q8CF90), rat (D3ZNT6). Other species not tested yet.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

Maf (musculoaponeurotic fibrosarcoma) family proteins belong to the basic-leucine zipper family of transcription factors and are divided into large Maf and small Maf proteins depending on their molecular size. Four types of large Maf transcription factors have been identified in mice and humans: MafA, MafB, c-Maf and NRL (1). MafA is a β-cell-specific transcription factor in adult pancreatic islets and regulates glucose-stimulated insulin secretion (2). MafB, on the other hand, is specifically expressed in glucagon-expressing α-cells in the pancreas of adult mice. In the embryonic or neonatal pancreas, immature β-cells first express MafB, whereas MafA is expressed in mature β-cells and is essential for the maintenance of the mature phenotype (3). Several studies have shown, that the expression of MafA is impaired in β-cells of rodents and humans with diabetes (3). In addition to its role in the pancreas, MafA is also involved in the development of neuronal cells. For example, MafA is involved in the development of glutamatergic neurons in the olfactory bulb (4) and of dorsal root ganglia in the spinal cord (5).

## Selected General References

Functional analysis of large MAF transcription factors and elucidation of their relationships with human diseases. Takahashi S et al. *Exp Anim* (2021) PubMed:33762508

Role of the Transcription Factor MAFA in the Maintenance of Pancreatic β-Cells. Nishimura W et al. *Int J Mol Sci* (2022) PubMed:35562869

Expression of Maf family proteins in glutamatergic neurons of the mouse olfactory bulb. Ito A et al. *Dev Neurobiol* (2022) PubMed:34679244

c-Maf is required for the development of dorsal horn laminae III/IV neurons and mechanoreceptive DRG axon projections. Hu J et al. *J Neurosci* (2012) PubMed:22514301

MafA is a glucose-regulated and pancreatic beta-cell-specific transcriptional activator for the insulin gene. Kataoka K et al. *J Biol Chem* (2002) PubMed:12368292

Access the online factsheet including applicable protocols at <https://susy-histosure.com/product/HS-521003> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.